



# TUFF TAQ™ DNA POLYMERASE

Reliable Activity Even After 3  
Months At Room Temperature



## KEY FEATURES:

99% active after 1 month at  
room temperature

>80% active after 3 months at  
room temperature

Ships at ambient temperature -  
No dry ice

Freeze/thaw up to 50 times

**TUFF Taq™ DNA Polymerase** is engineered with a proprietary peptide tag that confers extreme stability at room temperature. Robust, dependable and ideal for use in everyday PCR applications. Has 5' to 3' polymerase & exonuclease activity but no 3' to 5' exonuclease activity.

## KIT CONTENTS

- TUFF Taq™ DNA Polymerase
- 10X PCR buffer C: 0.8M Tris-HCL + 0.2M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> , 0.2% w/v Tween-20 (contains no Mg<sup>2+</sup> )
- 10X PCR buffer D: 0.8M Tris-HCL + 0.2M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + detergent (contains no Mg<sup>2+</sup> or detergent )
- 10x Solution E (Use with difficult templates eg GC rich or when non specific amplifications occur). Determine optimal working concentration empirically)
- 25mM MgCl<sub>2</sub>

## APPLICATIONS

- Ideal for most PCR Assays
- TA Cloning

## SHIPPING & STORAGE CONDITIONS

Shipping and temporary storage for up to 1 month at room temperature has no significant detrimental performance effects

**Routine storage:** -20°C

**Safety warnings:** Products and components should be handled only by persons trained in laboratory techniques. Wear suitable protective clothing such as laboratory overalls, gloves and safety glasses. Avoid contact with eyes and skin and wash immediately with water should contact occur.

**Usage restrictions:** For Research Use Only. Not for any animal or human therapeutic or diagnostic use. Some potential applications may require a license which is not provided by purchase of this product and users should obtain the license if required. Rebel BioScience products may not be re-sold, transferred, modified or used to manufacture commercial products without written approval from Rebel BioScience, LLC.

## RECOMMENDED PCR REACTION MIX FOR 100 µl

Component	Volume	Final Conc
TUFF Taq™ DNA Polymerase 5 U / µl	4µl	0.02 – 0.05 U / µl
10X PCR Buffer (A or B)	10 µl	1X
25mM MgCl <sub>2</sub>	6-10 µl	1.5-2.5mM
20mM dNTP mix	1 µl	200 µM
Forward Primer (10 pmol / µl)	1-3 µl	0.1-0.3 µM
Reverse Primer (10 pmol / µl)	1-3 µl	0.1-0.3 µM
DNA Template	5-20 µl	5-100 ng / µl
10X Solution E (if required, not for standard PCR)	0, 10, 20 or 30 µl	1X, 2X or 3X
PCR Grade H <sub>2</sub> O	Up to 100 µl	
<b>Total</b>	<b>100 µl</b>	

## RECOMMENDED PCR CYCLING CONDITIONS

Cycle Step	Temperature	Time	# Cycles
<b>Initial denaturation</b>	<b>95°C</b>	<b>3-5 min</b>	<b>1</b>
Denaturation	95°C	30-60 sec	26-35
Annealing	50-68°C	30-60 sec	
Elongation	72°C	1-4 min	
Final Elongation	72°C	5-10 min	1